

# The protective and therapeutic effects of Persian Gulf sea cucumber (*Holothuria leucospilota*) on Carbon tetrachloride - induced hepatotoxicity in Molly Fish (*Poecilia sphenops*)

Mohammadi A.<sup>1</sup>; Hosseinzadeh Sahaifi H.<sup>2\*</sup>; Naji T.<sup>3</sup>

Received: March 2019

Accepted: June 2019

## Abstract

Carbon tetrachloride (CCl<sub>4</sub>) is a well-known toxicant in exposed animals. It is assumed sea cucumber has medical properties. So, the protective and therapeutic effects of ethanol extract of the Persian Gulf sea cucumber (EEPGSC) on CCl<sub>4</sub>-induced hepatotoxicity in Molly Fish (*Poecilia sphenops*). A total 90 Molly fish were randomly divided into 9 experimental groups as follows: Group (C<sub>0</sub>): had no received injections. Group (C<sub>1</sub>): fish injected with olive oil (1 mL/kg, i.m.). Group (C<sub>2</sub>): injected with ethanol (100 mg/kg, i.m.) Group (C<sub>3</sub>): injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 1<sup>st</sup>. Group (T<sub>1</sub>): fish injected with CCl<sub>4</sub> (1 mL/kg, i.m.) and then with EEGPSC (100 mg/kg, i.m.). Group (T<sub>2</sub>): injected with CCl<sub>4</sub> (1 mL/kg, i.m.) then with EEGPSC (200 mg/kg, i.m.). Group (C<sub>4</sub>): had no received injections for 20 days, then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21. Group (T<sub>3</sub>): injected with EEGPSC (100 mg/kg, i.m.) for 20 days (given on alternate days) then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21. Group (T<sub>4</sub>): fish injected with EEGPSC (200 mg/kg, i.m.) for 20 days (given on alternate days) then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21. After the last injection, body length (cm), weight (g), GIS index and liver alkaline phosphatase (ALP), alanine aminotransferase (ALT) and Aspartate aminotransferase (AST) were determined. No significant difference detected in body weight and length compared to the control groups ( $p > 0.05$ ). Injection of the different levels of the EEGPSC (100 and 200 mg/kg, i.m.) decreased CCl<sub>4</sub>-induced ALP, ALT and AST levels compared to the C<sub>4</sub> groups ( $p < 0.05$ ). These results suggested EEGPSC had protective and therapeutic effect against CCl<sub>4</sub>-induced hepatotoxicity in Molly fish.

**Keywords:** Persian Gulf sea cucumber, Hepatotoxicity, Molly Fish

1-Pharmacy student in Department of Basic Sciences, Faculty of Pharmacy, Pharmaceutical Sciences Branch, Islamic Azad University (IAUPS), Tehran, Iran.

2-Associate of professor in Iranian Fisheries Science Research Institute, Agricultural Research, Education and Extension Organization, Tehran, Iran.

3- Department of Basic Sciences, Faculty of pharmacy and pharmaceutical sciences, Tehran Medical Science, Islamic Azad University, Tehran, Iran

Corresponding author's Email: h\_hosseinzadeh@yahoo.com

## Introduction

Liver is the organ which expose to several toxins and infections through hepatic vein in the body. Xenobiotic-metabolizing enzymes (mainly cytochrome P-450) are present in the liver where detoxify and eliminate toxicants. The toxicology of the liver is complicated by the diversity of liver injuries and the dissimilar mechanism through which the injuries are involved (Ujah *et al.*, 2013). Carbon tetrachloride (CCl<sub>4</sub>) is an experimental candidate toxicant extensively used to induce lipid peroxidation and toxicity and diseased state in animals (Peng *et al.*, 2010; Abdel Salam *et al.*, 2012 a, b; 2013 a, b). It is reported that CCl<sub>4</sub> promotes hepatocellular carcinoma incidence in rats ( ). There are varieties of enzymes include AST, ALT, ALP which elevated their serum concentrations are a powerful marker for liver damage (Abdel Salam *et al.*, 2012c). Recently, besides physicians, patient's interest increased to the use of botanical medicine and herbal remedies in liver disease management (Fogden and Neuberger, 2003; Karimi *et al.*, 2006; Hayatghaybi and Karimi, 2007; Karimi *et al.*, 2007; Saberivand *et al.*, 2010; Yousofi *et al.*, 2011). On the other hand there is growing interest for therapeutic drugs, products from marine sources (Esmat *et al.*, 2013).

*Holothuroidea* or sea cucumbers are abundant worm-like and soft-bodied echinoderms found in nearly every marine environment (Haider *et al.*, 2015). Sea cucumbers are the aquatic

creatures and have nutritional and therapeutic properties on human health (Mamelona *et al.*, 2007). Sea cucumbers are considered as a culinary delicacy and as a traditional cure for many illnesses in many Asian countries (Liu *et al.*, 2012). Some compounds isolated to date exhibit antimicrobial and anti-inflammatory, antioxidant, anticoagulant, anti-hyperlipidemic and immune modulatory (Hu *et al.*, 2012). It is reported the sea cucumber extract contains physiologically active phenolic compounds with antioxidant activity, which afforded a potential hepatoprotective activity against thioacetamide induced liver injury in a rat model (Esmat *et al.*, 2013). Recently, Dakrory *et al.* (2015) revealed *Holothuria atra* extract exhibits good hepatoprotective, curative and antioxidant potential against DMBA-induced hepato-renal dysfunction in rats. The therapeutic use of sea cucumbers for healing is established, where they used for joint pain, tendonitis and sprains. For instance, sea cucumbers extract decrease serum aminotransferases (ASAT and ALAT) and ALP as well as malondialdehyde (MDA) while increasing the serum albumin, glutathione reduced (GSH) levels in bile duct ligation in rats (Fahmy, 2015). Also, *Holothuria arenicola* extract normalized the antioxidant enzyme, glutathione-S-transferase (GST), superoxide dismutase (SOD) and catalase (CAT) activities (Fahmy, 2015).

Based on the literature, scarce information exists on medical properties of the Persian Gulf sea cucumber (*Holothuria leucospilota*). In a sole report, Delghandi Moghadamet *al.* (2016) studied effect of Persian Gulf sea cucumber (*Holothuria leucospilota*) on maturation of mice. So, the protective and therapeutic effects of ethanol extract of EEPGSC on CCl<sub>4</sub>-induced hepatotoxicity in Molly Fish (*Poecilia sphenops*) were studied.

## Material and methods

### *Sample preparation*

Samples of Persian Gulf sea cucumber (*Holothuria leucospilota*) were collected from Persian Gulf, at the depth of 25-30 meters in 2016. Then samples transported and stored at -20°C in laboratory of Fisheries, Department of Pharmacology, Islamic Azad University, Tehran, Iran.

### *Persian Gulf sea cucumber extract*

The Persian Gulf sea cucumber (EEPGSC) (2000 g wet weight) was cut into small pieces (1 cm), they were put in freeze dried until the tissue was dry (150 g dry weight). Extracts of powder sea cucumber were obtained by using three different solvents including: n-hexane, diethyl ether and methanol. After 24 hours of exposure in n-hexane, the extract was concentrated under low pressure at 30°C by rotary evaporation. The diethyl ether extract was ready after 48 hours, and then the solvent was removed by rotary evaporation at 35°C. The methanol extract was ready after 72

hours, then the solvent was removed by rotary evaporation at 40°C. Ether-methanol was formed by adding Ether in order to separate the methanol-aqueous extract, then the upper phase was separated by separating funnel. The upper phase was combined by n-butanol and aqueous extract was separated by separating funnel. Each extract was shaken by mechanical shaking at room temperature (25°C). All processes were carried out on dark condition. Finally, both crude extracts were kept in freezer (Bligh *et al.*, 1959).

### *Experimental groups*

A total 90 Molly fish (*Poecilia sphenops*) (3±0.2) were randomly divided into 9 experimental groups (n=10). The fish were kept in 2 m<sup>3</sup> tanks with a flow-through circuit, suitable aeration and filtration system and natural photoperiod. The water temperature ranged from 25.1 to 27.8°C. The environmental parameters, mortality and food intake were recorded daily. Before the initiation of the study, fish kept for a week for adaptation to the experimental condition, and then randomly divided into experimental groups as follow in Table 1.

### *Serum Biomarkers for Liver Functions Tests*

Then 24 h after the last injection, fish from each group randomly selected, euthanized and biometric indexes body length (cm), weight (g), and GIS index were determined.

**Table 1: the experimental groups.**

Group (C <sub>0</sub> )	Had no received injections and fish kept as control.
Group (C <sub>1</sub> )	Fish injected with olive oil (1 ml/kg, i.m.) for 20 days (given on alternate days).
Group (C <sub>2</sub> )	Fish injected with ethanol (100 mg/kg, i.m.) for 20 days (given on alternate days).
Group (C <sub>3</sub> )	Fish injected with CCl <sub>4</sub> (1 ml/kg, i.m.) on day 1 <sup>st</sup> .
Group (T <sub>1</sub> )	Fish injected with CCl <sub>4</sub> (1 ml/kg, i.m.) and then injected with EEGGSC (100 mg/kg, i.m.) for 20 days (given on alternate days).
Group (T <sub>2</sub> )	Fish injected with CCl <sub>4</sub> (1 ml/kg, i.m.) and then injected with EEGGSC (200 mg/kg, i.m.) for 20 days (given on alternate days).
Group (C <sub>4</sub> )	Had no received injections for 20 days, then injected with CCl <sub>4</sub> (1 ml/kg, i.m.) on day 21.
Group (T <sub>3</sub> )	Fish injected with EEGGSC (100 mg/kg, i.m.) for 20 days (given on alternate days) then injected with CCl <sub>4</sub> (1 ml/kg, i.m.) on day 21.
Group (T <sub>4</sub> )	Fish injected with EEGGSC (200 mg/kg, i.m.) for 20 days (given on alternate days) then injected with CCl <sub>4</sub> (1 ml/kg, i.m.) on day 21.

Also, the liver samples obtained and the tissue alkaline phosphatase (ALP), alanine aminotransferase (ALT) and Aspartate aminotransferase (AST) were determined using commercial detecting kits (Reitman and Frankel, 1957; Belfield and Goldberg, 1971).

#### Statistical analyses

Effect of EEGGSC was analyzed by repeated measure two-way analysis of variance (ANOVA) using SPSS 21.0 for Windows (SPSS, Inc., Chicago, IL, USA). Data is presented as mean±SEM. For treatment showing a main effect by ANOVA, means compared by Tukey-Kramer test.  $p < 0.05$  was considered as significant differences between treatments.

#### Results


The protective and therapeutic effects of EEGGSC on CCl<sub>4</sub>-induced hepatotoxicity in Molly Fish (*Poecilia sphenops*) is presented in Figs. 1-6. According to the results despite there was difference between the experimental groups, no significant difference observed in body weight gain

compared to the control groups ( $p > 0.05$ ) (Fig. 1).


The protective and therapeutic effects of EEGGSC on body length (cm) of Molly fish was presented in Fig. 2. According to the results, no significant difference detected on body length (cm) compared to the control group ( $p > 0.05$ ), however, the mean body length was higher in fish injected with olive oil (1 mL/kg, i.m.) for 20 days (given on alternate days).

As seen in Fig. 3, no significant difference detected in GIS index in fish injected with EEGGSC, olive oil, ethanol and CCl<sub>4</sub> compared to the control group ( $p > 0.05$ ).


As seen in Fig. 4, no significant difference detected in tissue ALP levels in fish injected with olive oil (1 mL/kg, i.m.) for 20 days (given on alternate days) and fish injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 1<sup>st</sup> compared to the control group ( $p > 0.05$ ). Furthermore, tissue ALP level was significantly higher in fish injected with CCl<sub>4</sub> compared to the compared to the control group ( $p < 0.05$ ).




**Figure 1:** The protective and therapeutic effects of EEGSC on body weight (g) of Molly fish (*Poecilia sphenops*). EEGSC: ethanol extract of the Persian Gulf sea cucumber; C<sub>0</sub>: had no received injections and fish kept as control; C<sub>1</sub>: fish injected with olive oil (1 mL/kg, i.m.) for 20 days (given on alternate days); C<sub>2</sub>: fish injected with ethanol (100 mg/kg, i.m.) for 20 days (given on alternate days); C<sub>3</sub>: fish injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 1<sup>st</sup>; T<sub>1</sub>: fish injected with CCl<sub>4</sub> (1 mL/kg, i.m.) and then injected with EEGSC (100 mg/kg, i.m.) for 20 days (given on alternate days); T<sub>2</sub>: fish injected with CCl<sub>4</sub> (1 mL/kg, i.m.) and then injected with EEGSC (200 mg/kg, i.m.) for 20 days (given on alternate days); C<sub>4</sub>: had no received injections for 20 days, then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21; T<sub>3</sub>: fish injected with EEGSC (100 mg/kg, i.m.) for 20 days (given on alternate days) then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21; T<sub>4</sub>: fish injected with EEGSC (200 mg/kg, i.m.) for 20 days (given on alternate days) then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21.



**Figure 2:** The protective and therapeutic effects of EEGSC on body length (cm) of Molly fish (*Poecilia sphenops*). EEGSC: ethanol extract of the Persian Gulf sea cucumber; C<sub>0</sub>: had no received injections and fish kept as control; C<sub>1</sub>: fish injected with olive oil (1 mL/kg, i.m.) for 20 days (given on alternate days); C<sub>2</sub>: fish injected with ethanol (100 mg/kg, i.m.) for 20 days (given on alternate days); C<sub>3</sub>: fish injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 1<sup>st</sup>; T<sub>1</sub>: fish injected with CCl<sub>4</sub> (1 mL/kg, i.m.) and then injected with EEGSC (100 mg/kg, i.m.) for 20 days (given on alternate days); T<sub>2</sub>: fish injected with CCl<sub>4</sub> (1 mL/kg, i.m.) and then injected with EEGSC (200 mg/kg, i.m.) for 20 days (given on alternate days); C<sub>4</sub>: had no received injections for 20 days, then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21; T<sub>3</sub>: fish injected with EEGSC (100 mg/kg, i.m.) for 20 days (given on alternate days) then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21; T<sub>4</sub>: fish injected with EEGSC (200 mg/kg, i.m.) for 20 days (given on alternate days) then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21.



**Figure 3:** The protective and therapeutic effects of EEPGSC on GSI index in Molly fish (*Poecilia sphenops*). EEPGSC: ethanol extract of the Persian Gulf sea cucumber; C<sub>0</sub>: had no received injections and fish kept as control; C<sub>1</sub>: fish injected with olive oil (1 mL/kg, i.m.) for 20 days (given on alternate days); C<sub>2</sub>: fish injected with ethanol (100 mg/kg, i.m.) for 20 days (given on alternate days); C<sub>3</sub>: fish injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 1<sup>st</sup>; T<sub>1</sub>: fish injected with CCl<sub>4</sub> (1 mL/kg, i.m.) and then injected with EEPGSC (100 mg/kg, i.m.) for 20 days (given on alternate days); T<sub>2</sub>: fish injected with CCl<sub>4</sub> (1 mL/kg, i.m.) and then injected with EEPGSC (200 mg/kg, i.m.) for 20 days (given on alternate days); C<sub>4</sub>: had no received injections for 20 days, then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21; T<sub>3</sub>: fish injected with EEPGSC (100 mg/kg, i.m.) for 20 days (given on alternate days) then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21; T<sub>4</sub>: fish injected with EEPGSC (200 mg/kg, i.m.) for 20 days (given on alternate days) then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21.




**Figure 4:** The protective and therapeutic effects of EEPGSC on liver ALP levels in Molly fish (*Poecilia sphenops*). EEPGSC: ethanol extract of the Persian Gulf sea cucumber; ALP: alkaline phosphatase; C<sub>0</sub>: had no received injections and fish kept as control; C<sub>1</sub>: fish injected with olive oil (1 mL/kg, i.m.) for 20 days (given on alternate days); C<sub>2</sub>: fish injected with ethanol (100 mg/kg, i.m.) for 20 days (given on alternate days); C<sub>3</sub>: fish injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 1<sup>st</sup>; T<sub>1</sub>: fish injected with CCl<sub>4</sub> (1 mL/kg, i.m.) and then injected with EEPGSC (100 mg/kg, i.m.) for 20 days (given on alternate days); T<sub>2</sub>: fish injected with CCl<sub>4</sub> (1 mL/kg, i.m.) and then injected with EEPGSC (200 mg/kg, i.m.) for 20 days (given on alternate days); C<sub>4</sub>: had no received injections for 20 days, then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21; T<sub>3</sub>: fish injected with EEPGSC (100 mg/kg, i.m.) for 20 days (given on alternate days) then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21; T<sub>4</sub>: fish injected with EEPGSC (200 mg/kg, i.m.) for 20 days (given on alternate days) then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21. There are significant differences between groups with different superscripts in a column ( $p < 0.05$ ).

Injection of the different levels of the EEGGSC (100 and 200 mg/kg, i.m.) significantly decreased CCl<sub>4</sub>-induced ALP levels compared to the C<sub>4</sub> groups ( $p < 0.05$ ). These results suggest EEGGSC has protective and therapeutic effect against CCl<sub>4</sub>-induced hepatotoxicity in Molly fish.

No significant difference observed in liver ALT levels in Molly fish injected with olive oil (1 mL/kg, i.m.) for 20 days and ethanol (100 mg/kg, i.m.) for 20 days compared to control group ( $p > 0.05$ ) but injection of the CCl<sub>4</sub> (1 mL/kg) on day 1<sup>st</sup> significantly

increased liver ALT levels in comparison to control group ( $p < 0.05$ ). Additionally, liver ALT levels significantly increased in group C<sub>4</sub> (no received injections for 20 days, then injected with 1 mL/kg of CCl<sub>4</sub> on day 21) compared to control group) while injection of the EEGGSC (100 and 200 mg/kg, i.m.) for 20 days significantly decreased CCl<sub>4</sub>-induced ALT levels ( $p < 0.05$ ) Fig. 5). These results suggest EEGGSC had protective and therapeutic effect CCl<sub>4</sub>-induced hepatotoxicity in Molly fish.




**Figure 5:** The protective and therapeutic effects of EEGGSC on liver ALT levels in Molly fish (*Poecilia sphenops*). EEGGSC: ethanol extract of the Persian Gulf sea cucumber; ALT: alanine aminotransferase; C<sub>0</sub>: had no received injections and fish kept as control; C<sub>1</sub>: fish injected with olive oil (1 mL/kg, i.m.) for 20 days (given on alternate days); C<sub>2</sub>: fish injected with ethanol (100 mg/kg, i.m.) for 20 days (given on alternate days); C<sub>3</sub>: fish injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 1<sup>st</sup>; T<sub>1</sub>: fish injected with CCl<sub>4</sub> (1 mL/kg, i.m.) and then injected with EEGGSC (100 mg/kg, i.m.) for 20 days (given on alternate days); T<sub>2</sub>: fish injected with CCl<sub>4</sub> (1 mL/kg, i.m.) and then injected with EEGGSC (200 mg/kg, i.m.) for 20 days (given on alternate days); C<sub>4</sub>: had no received injections for 20 days, then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21; T<sub>3</sub>: fish injected with EEGGSC (100 mg/kg, i.m.) for 20 days (given on alternate days) then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21; T<sub>4</sub>: fish injected with EEGGSC (200 mg/kg, i.m.) for 20 days (given on alternate days) then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21. There are significant differences between groups with different superscripts in a column ( $p < 0.05$ ).

As seen in Fig. 6, CCl<sub>4</sub> (1 mL/kg, i.m.) on day 1<sup>st</sup> significantly elevated tissue

AST level compared to the control group ( $p < 0.05$ ). Also, tissue AST levels

significantly decreased in fish injected with CCl<sub>4</sub> (1 mL/kg, i.m.) and then injected with EEPGSC (100 mg/kg, i.m.) for 20 days (given on alternate days) ( $p < 0.05$ ). Injection of the different levels of the EEPGSC (200 mg/kg, i.m.) for 20 days (given on

alternate days) then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21 significantly decreased CCl<sub>4</sub>-induced AST compared to the C<sub>4</sub> group ( $p < 0.05$ ). These results suggest EEPGSC was able to minimize CCl<sub>4</sub>-induced hepatotoxicity in Molly fish.



**Figure 6:** The protective and therapeutic effects of EEPGSC on liver AST levels in Molly fish (*Poecilia sphenops*). EEPGSC: ethanol extract of the Persian Gulf sea cucumber; AST: Aspartate aminotransferase; C<sub>0</sub>: had no received injections and fish kept as control; C<sub>1</sub>: fish injected with olive oil (1 mL/kg, i.m.) for 20 days (given on alternate days); C<sub>2</sub>: fish injected with ethanol (100 mg/kg, i.m.) for 20 days (given on alternate days); C<sub>3</sub>: fish injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 1<sup>st</sup>; T<sub>1</sub>: fish injected with CCl<sub>4</sub> (1 mL/kg, i.m.) and then injected with EEPGSC (100 mg/kg, i.m.) for 20 days (given on alternate days); T<sub>2</sub>: fish injected with CCl<sub>4</sub> (1 mL/kg, i.m.) and then injected with EEPGSC (200 mg/kg, i.m.) for 20 days (given on alternate days); C<sub>4</sub>: had no received injections for 20 days, then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21; T<sub>3</sub>: fish injected with EEPGSC (100 mg/kg, i.m.) for 20 days (given on alternate days) then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21; T<sub>4</sub>: fish injected with EEPGSC (200 mg/kg, i.m.) for 20 days (given on alternate days) then injected with CCl<sub>4</sub> (1 mL/kg, i.m.) on day 21. There are significant differences between groups with different superscripts in a column ( $p < 0.05$ ).

## Discussion

To the best of this study is first report on the protective and therapeutic effects of Persian Gulf sea cucumber (*Holothuria leucospilota*) on Carbon tetrachloride -induced hepatotoxicity in Molly fish. As detected, no significant difference detected in body weight gain and body length but different levels of the EEPGSC (100 and 200 mg/kg, i.m.)

significantly decreased CCl<sub>4</sub>-induced ALP, ALT and AST levels.

CCl<sub>4</sub> as an applicable candidate for other halogenated hydrocarbons has been one of the most broadly studied hepatotoxicant to date (Weber *et al.*, 2003). Acute poisoning with CCl<sub>4</sub> becomes manifested as a multi-organ disorder (Gitlin, 1996). It seems that liver is very vulnerable to this messy



hepatotoxin and a single exposure can quickly lead to severe necrosis and steatosis. Organ function tests are useful to evaluate amount or severity of CCl<sub>4</sub>-induced toxicity. In this line, assaying activities of several enzyme like LDH, AST, ALT, CPK, ALP, Glutathione-S-transferase (GST) and Glucuronyltransferase, catalase, superoxide dismutase, glutathione peroxidase has been frequently used to determine CCl<sub>4</sub>-induced toxicity (Peng *et al.*, 2010; Weerachayaphorn *et al.*, 2010; Abdel Salam *et al.*, 2012 a, b; 2013 a, b). Other studies that have a mechanistic view regarding CCl<sub>4</sub>-induced toxicity usually measure biomarker of oxidative stress like reduced glutathione (GSH) and malondialdehyde (MDA) (Wills *et al.*, 2006).

It is reported *Holothuria atra* extract decrease serum AST, ALT and ALP as well as MDA levels in bile duct ligation in rats (Fahmy, 2015). Also, the sea cucumber extracts increase GSH and normalized GST, SOD, CAT levels (Fahmy, 2015). So, it seems sea cucumber extracts have potential role against hepatotoxicity in rat model. In the current study, because of the limitations we were not able to determine effects of the EEPGSC on reactive oxidation species (ROS) like MDA, SOD and CAT. The modern medicinal system relies heavily on synthetic chemicals being used as drugs, but these unnatural synthetic drugs often pose serious side effects. Therefore, the development of novel

chemotherapeutic agents would play a key role in the treatment of many refractory diseases.

The Persian Gulf sea cucumber (*Holothuria leucospilota*) is a marine invertebrate of the phylum Echinoderm and the class *Holothuroidea* found on the sea floor worldwide (Farjami *et al.*, 2014). Dry sea cucumber contains approximately 20 mg/g glucosylceramide. It is reported dietary sea cucumber containing glucosylceramide decreased liver cholesterol and triglyceride compared to glucosylceramide free sea cucumber in mice but not affect body weight (Hossain *et al.*, 2011). Polysaccharides can act as stimulators of bile acid synthesis and circulation and increasing the fecal excretion of bile acids so that fewer bile acids return to the liver (Hossain *et al.*, 2011). The present study explores the protective and curative roles of the EEPGSC on ALP, ALT and AST which the findings were similar to Fahmy (2015). As observed in the current study, injection of CCl<sub>4</sub> elevated ALT, AST and ALP in Molly fish. Increase in ALP levels confirmed the damage produced by the bile duct ligation in rats (Baldo *et al.*, 2011). This increase may be attributable to the retention of bile salts that damaged the membrane and consequently leads to the passing of the ALP enzyme into circulation (Baldo *et al.*, 2012). In this regard, Esmat *et al.* (2013), reported administration of the *Holothuria leucospilota* extract attenuated the hepatic enzymes as evident by 77, 70

and 47% improvement in the activities of serum AST, ALT and ALP respectively in the rats. In the current study, our hypothesis for using fish as animal model firstly was because of anatomical and comparative physiological difference among animal (marine vs. mammalian) and secondly for determining possible effects EEPGSC in fish as sea food source for human. These results suggested sea cucumber not only could use as medical substance but also, it can be used as food additive in fishery industry to improve quality of the fish meat and its safety for human consumption (Farjami *et al.*, 2014).

In conclusion these results suggested EEPGSC has protective and therapeutic effect against CCl<sub>4</sub>-induced hepatotoxicity in Molly fish. To date, several bioactive components and medical properties of the sea cucumber have been identified. The obtained results of the current study could count as new medical properties of the Persian Gulf sea cucumber in medicine, pharmacology and even veterinary. However, further researches needed to determine accuracy of the results and the possible molecular and cellular mechanisms for effect of EEPGSC for clinical trials.

### Acknowledgement

Authors would like to thanks to Professor Motalebi for financial supporting and Dr. Nazemi for field assistance of the project.

### References

- Abdel Salam, O.M.E., Sleem, A.A. and Shafee, N., 2012a.** Effect of *Crataegus extract* on carbon tetrachloride-induced hepatic damage. *Comparative Clinical Pathology*, 21, 1719-1726.
- Abdel Salam, O.M.E, Sleem, A.A. and Shafee, N., 2012b.** *Hypericum perforatum* protects against hepatic injury induced by carbon tetrachloride. *Comparative Clinical Pathology*, 21, 1149-1157
- Abdel Salam, O.M.E, Sleem, A.A. and Shafee, N., 2013a.** The dopamine agonist *piribedil* exerts hepatoprotective effects on carbon tetrachloride-induced hepatic damage. *Comparative Clinical Pathology*, 22, 413-419.
- Abdel Salam, O.M.E., Sleem, A.A. and Omara, E., 2013b.** Micronised purified *flavonoid* fraction alleviates the carbon tetrachloride-induced hepatic injury. *Comparative Clinical Pathology*, 22, 1145-1154.
- Abdel Salam, O.M.E., Sleem, A.A. and Shafee, N., 2012c.** Hepatoprotective effects of *Cynara* extract and silymarin on carbon tetrachloride-induced hepatic damage in rats. *Comparative Clinical Pathology*, DOI 10.1007/s00580-012-1675-3.
- Baldo, G., Kretzmann, N.A., Tieppo, J., Filho, G.P., Cruz, C.U., Meurer, L., Da Silveira, T.R., dos Santos, J.L., Marroni, C.A., Marroni, N.P., Giugliani, R. and Matte, U.,**

2011. Bone marrow cells reduce collagen deposition in the rat model of common bile duct ligation. *Journal of Cell Science and Therapy*, 2, 1-12.
- Belfield, A. and Goldberg, D.M., 1971.** Revised assay for serumphenyl phosphatase activity using 4-amino-antipyrine. *Enzyme*, 12(5), 561–573.
- Bligh, E.G. and Dyer, W.J., 1959.** A rapid method for total lipid extraction and purification. *Canadian Journal of Biochemistry and Physiology*, 37, 911-7.
- Dakrory, A.I., Fahmy, S.R., Soliman, A.M., Mohamed, A.S. and Amer, S.A.M., 2015.** Protective and curative effects of the sea cucumber *Holothuria atra* extract against DMBA-induced hepatorenal diseases in rats. *BioMed Research International*.  
<http://dx.doi.org/10.1155/2015/5636>  
52
- Delghandi Moghadam, F., Baharara, J., Zafar Balanezhad, S., Jalali, M. and Amini, E., 2016.** Effect of *Holothuria leucospilota* extracted saponin on maturation of mice oocyte and granulosa cells. *Research in Pharmaceutical Sciences*, 11(2), 130-137.
- Esmat, A.Y., Said, M.M., Soliman, A.A., El-Masry, K.S.H. and Badiea, E.A., 2013.** Bioactive compounds, antioxidant potential, and hepatoprotective activity of sea cucumber (*Holothuria atra*) against thioacetamide intoxication in rats. *Nutrition*, 29(1), 258–267.
- Fahmy, S.R., 2015.** Anti-fibrotic effect of *Holothuria arenicola* extract against bile duct ligation in rats. *Comp. Altern. Med.* 15, 14. DOI 10.1186/s12906-015-0533-7
- Farjami, B., Nematollahi, M., Moradi, Y. and Nazemi, M., 2014.** Derivation of extracts from Persian Gulf sea cucumber (*Holothuria leucospilota*) and assessment of its antifungal effect. *Iranian Journal of Fisheries Sciences*, 13(4), 785-795.
- Fogden, E. and Neuberger, J., 2003.** Alternative medicines and the liver. *Liver International*, 23, 213-220
- Gitlin, N., 1996.** Clinical aspects of liver disease caused by industrial and environmental toxins. In: Zakim D, Boyer TD, editors. *Hepatology. A Textbook of Liver Disease*. Philadelphia, USA: WB Saunders, pp. 1018-1050.
- Haider, M.S., Sultana, R., Jamil, K., Lakht, Z., Tarar, O.M., Shirin, K. and Afzal, W., 2015.** A Study on proximate composition, amino acid profile, fatty acid profile and some mineral contents in two species of sea cucumber. *Journal of Animal and Plant Sciences*, 25(1), 168-175.
- Hayatghaybi, H. and Karimi, I., 2007.** Hypercholesterolemia effect of drug-type *Cannabis sativa L.* seed (marijuana seed) in guinea pig. *Pakistan Journal of Nutrition*, 6, 56-59.
- Hossain, Z., Sugawara, T., Aida, K., Hirata, T., 2011.** Effect of dietary

- glucosylceramide from sea cucumber on plasma and liver lipids in cholesterol-fed mice. *Fisheries Science*, 77, 1081–1085.
- Hu, XQ. Xu, J., Xue, Y., Li, Z.J., Wang, J.F., Wang, J.H., Xue, C.H. and Wang, Y.M., 2012.** Effects of bioactive components of sea cucumber on the serum, liver lipid profile and lipid absorption. *Bioscience, Biotechnology, and Biochemistry*, 76 (12), 2214–2218.
- Karimi, I. and Hayatghaibi, H., 2006.** Effect of *Cannabis Sativa L. seed* (hempseed) on serum lipid and protein profiles of rat. *Pakistan Journal of Nutrition*, 5, 585–588.
- Karimi, I., Hayatghaibi, H., Yousofi, J., Saberivand, A. and Zavareh, S., 2007.** Effects of nondrug type *Cannabis sativa L.* (hempseed) on hematological parameters in guinea pig. *J. Cardiova Hemato Dis -Drug Targets*, 7, 288–290.
- Liu, X., Sun, Z., Zhang, M., Meng, X., Xia, X., Yuan, W., Xue, F. and Liu, C., 2012.** Antioxidant and antihyperlipidemic activities of polysaccharides from sea cucumber *Apostichopus japonicas*. *Carbohydrate Polymers*, 90, 1664–1670.
- Mamelona, J., Pelletier, E., Girard-Lalancette, K., Legault, J., Karboune, S. and Kermasha, S., 2007.** Quantification of phenolic contents and antioxidant capacity of Atlantic sea cucumber, *Cucumaria frondosa*. *Food Chemistry*, 104, 1040–1047.
- Peng, W.H., Tien, Y.C., Huang, C.Y., Huang, T.H., Liao, J.C., Kuo, C.L and Lin, Y.C., 2010.** *Fraxinus rhynchophylla* ethanol extract attenuates carbon tetrachloride-induced liver fibrosis in rats via down-regulating the expressions of uPA, MMP-2, MMP-9 and TIMP-1. *Journal of Ethnopharmacology*, 127, 606–613.
- Reitman, S. and Frankel, S., 1957.** A colorimetric method for the determination of serum glutamic oxalacetic and glutamic pyruvic transaminases. *American Journal of Clinical Pathology*, 28(1), 56–63.
- Saberivand, A., Karimi, I., Becker, L.A., Moghaddam, A., Azizi-Mahmoodjigh, S., Yousefi, M. and Zavareh, S., 2010.** The effects of *Cannabis sativa L.* seed (hempseed) in the ovariectomized rat model of menopause. *Methods find Exp. Journal of Clinical Pharmacology*, 32, 467–473.
- Ujah, O.F., Ujah, I.R., Johnson, J.T., Ekam, V.S. and Udenze, E.C.C., 2013.** Hepatoprotective property of ethanolic leaf extract of *Moringa oleifera* on carbon tetrachloride (CCl<sub>4</sub>) induced hepatotoxicity. *Journal of Natural Product and Plant Resources*, 3 (2), 15–22.
- Weber, L.W.D., Boll, M. and Stampfl, A., 2003.** Hepatotoxicity and mechanism of action of haloalkanes: carbon tetrachloride as

a toxicological model.  
*Critical Reviews in Toxicology*, 33,  
105-136.

**Weerachayaphorn, J.,  
Chuncharunee, A., Jariyawat, S.,  
Lewchalermwong, B,  
Amonpatumrat, S., Suksamrarn,  
A. and Piyachaturawat, P., 2010.**  
Protection of centrilobular necrosis  
by *Curcuma comosa Roxb.* In carbon  
tetrachloride-induced mice liver  
injury. *Journal  
of Ethnopharmacology*, 129, 254-  
260.

**Wills, P.J. and Asha, V.V., 2006.**  
Protective effect of *Lygodium  
flexuosum* (L.) Sw. extract against  
carbon tetrachloride-induced acute  
liver injury in rats. *Journal  
of Ethnopharmacology*, 108, 320-  
326.

**Yousofi, M., Saberivand, A., Becker,  
L.A. and Karimi, I., 2011.** The  
effects of *Cannabis sativa L.* seed  
(hemp seed) on reproductive and  
neurobehavioral end points in rats.  
*Developmental Psychobiology*, 53,  
402-412.