



Mineral and elemental compositions of suspended particulate matter (SPM) and sediments from major rivers in the Southeastern Nigeria

Etuk H.S.¹; Ebong G.A.^{1*}; Ekot A.E.²; Ekpo B.O.³; Ayi A.A.⁴

Received: July 2023

Accepted: November 2023

Abstract

Mineral and elemental compositions of suspended particulate matter (SPM) and sediments from Great Kwa, Calabar, Cross River, Imo, and Qua Iboe Rivers in the Southeastern part of Nigeria were assessed in this study. SPM samples were obtained by the collection of water samples from the studied Rivers, passing the water through dried and weighed Whatman fiberglass filters and the sediments obtained were dried to a constant weight. Bottom sediments were obtained using Van Veen Grab sampler, transported to the laboratory, and air-dried at room temperature for seven days. These samples were subjected to Energy Dispersive X-Ray Analysis (EDX), X-ray diffraction analysis (XRD), and Scanning electron microscope (SEM). EDX analysis revealed that, high levels of Si and Al were common for all the studied samples. Silicate mineral was observed in all the sediments. XRD revealed that Quartz, clays, micas, hornblende, potassium-feldspars and mafics were the major minerals in the SPM and sediments. Quartz, gypsum and gibbsite minerals higher during the high than in low tide. Zircon, amphibole, pyroxene, biotite, pyrite, and iron hydroxide were identified by SEM in the samples. The mineral and elemental compositions of SPM and sediments from the studied rivers revealed and the pollution status established.

Keywords: Energy Dispersive X-Ray Analysis, X-ray diffraction analysis, Scanning electron microscope, Water pollution, Mineralogical study, Southeastern Nigeria

1-Department of Chemistry, University of Uyo, P.M.B. 1017, Uyo, Akwa Ibom State, Nigeria.

2-Department of Geology, Akwa Ibom State University, Mkpato Enin, P. M. B. 1167 Uyo, Nigeria

3-Exploration & Production Department Division, Epc Complex Site, Eleme, Port Harcourt, River State.

4-Department of Pure & Applied Chemistry, University Of Calabar, Calabar.

*Corresponding author's Email: godwinebong@uniuyo.edu.ng

Introduction

The quality of water available for utilization by human has direct effect on the health status (Lin *et al.*, 2022; Luvhimbi *et al.*, 2022). The quality of water in the Niger Delta area of Nigeria has been greatly affected by the anthropogenic factor especially the oil exploration and exploitation activities within the area (Ebong and Etuk, 2016; Ewim *et al.*, 2023; Iwegbue *et al.*, 2023). Reports have shown that prolonged human exposure to polluted water may result in water-related diseases such as dysentery, cholera, hepatitis, and diarrhea (Magana-Arachchi and Wanigatunge, 2020; Lin *et al.*, 2022). High levels of toxic substances may have resulted in the extinction of some aquatic organisms within the oil producing communities of Nigeria (Bebeteidoh *et al.*, 2020; Yinet *et al.*, 2022). The aquatic ecosystem has high tendency of being polluted since it receives contaminants from numerous sources including municipal discharges, industrial, agricultural, and domestic wastes (Akhtar *et al.*, 2021). Elevated levels of metals in aquatic ecosystem are risky to all biological lives including human and is an indication of anthropogenic impact on the water quality rather than geological enrichment (Ebong *et al.*, 2006; Algül and Beyhan, 2020; Sojka and Jaskuła, 2022). Water sediments accumulate high levels of toxic substances mostly metals and subsequently release them into the water channel (Ebong and Etuk, 2016; Bartoszek *et al.*, 2022). The natural geological process like weathering of

rocks also have negative impact on the quality of water and sediments in aquatic environment (Baba and Gündüz, 2017; Yoshimura *et al.*, 2021). SPM and sediments in the aquatic environment are affected by both the natural and human-induced factors and it has negative impacts on both human and aquatic lives (Baba and Gündüz, 2017; Shil and Singh, 2019). However, if the anthropogenic rate at which sediments are enriched in a water system is higher than the natural source, it could be an indication of pollution (Chen *et al.*, 2014).

SPM is also a fundamental component of geochemical, geological, and biological sequence in the aquatic ecosystem due to its abundance, physicochemical properties, and mobility (Fernandes *et al.*, 2019). SPM in aquatic environment consists of materials that do not sediment under some hydrochemical and hydrodynamic conditions and can be retained on a filter paper with porosity of 0.45 μm (Walch *et al.*, 2022). SPM links the pelagic and benthic zones together in aquatic environment by sedimentation and re-suspension (Sutherland *et al.*, 2022). In the aquatic ecosystem, SPM has the potential of absorbing dissolved trace metals because of its large surface area and reactivity (Li *et al.*, 2019). Reports have shown that SPM carries majority of the solid contaminants from the soil environment to the aquatic environment (Zeng *et al.*, 2019; Fan *et al.*, 2021; 2023).

Studies have shown that the major components of SPM are mica, carbonate, phyllosilicates, plagioclase,

quartz, potassium feldspar, and clay minerals as well as chlorite, kaolinite, montmorillonite, and illite (Sunet *et al.*, 2020; Kumari and Mohan, 2021). Clay minerals emanate mostly from the chemical and physical weathering of rocks and from the final product of soil genesis (Biscaye *et al.*, 1997; Kumari and Mohan, 2021). Clay minerals are divided into four main groups namely: kaolinite, chlorite, illite, and smectite (Han *et al.*, 2022). The kaolinite group is mainly found in the humid tropical climatic zones of South America, Africa, and Indonesia in ferrasols soils (Ouyang *et al.*, 2021). The chlorites originate from mechanical weathering of rocks and are easily degraded during soil genesis. The illite group originates from the temperate and cold climate areas, mainly from Podzolic soils. Mica minerals originate mainly from direct mechanical weathering of igneous and metamorphic rocks. These clay minerals and quartz constitute the principal minerals of loess deposits (Pye and Blott, 2006; Chmielowska and Salata, 2020). Feldspar and mafic (mainly amphibole and pyroxene) minerals are derived from mechanical weathering of igneous and metamorphic rocks (Kloprogge and Wood, 2020; Macheyeke *et al.*, 2020).

SPM also have some organic components such as microorganisms and organic particles including cellular debris, detritus, and extracellular polymers (Walch *et al.*, 2022). Hence, it can affect the turbidity, colour, nutrient availability, and microbial load, clarity and availability of oxygen in aquatic

ecosystem (Hope *et al.*, 2020; Lednicka *et al.*, 2022; Nair and Nayak, 2023). SPM controls eutrophication as well as the bioavailability and mobility of organic and inorganic substances in aquatic environment (Patrolecco *et al.*, 2010; Vidmar *et al.*, 2017; Gordon *et al.*, 2020). Recent reports have indicated that SPM influences the presence of some emerging contaminants such as nanoplastics, nanoparticles, soots etc in aquatic ecosystem (Wagner *et al.*, 2018; Li *et al.*, 2019; Surette and Nason, 2019).

Sediments in water emanate from SPM and weathering of rocks, sediments are made up of and are consist mainly of calcite, clay minerals (kaolinite, montmorillonite, and illite), feldspar, and quartz in a silicate form (Saito *et al.*, 2017; Tian *et al.*, 2021). The other minerals found in river sediment included plagioclase, pyroxenes, muscovite, K-feldspars, biotite, and hornblende (Kimeli *et al.*, 2021). However, the compositions of sediments depend solely on the mineral compositions of the mother rock. Silicates in sediments consist of mainly quartz, clays, micas, feldspars, and mafic minerals (Kimeli *et al.*, 2021). Quartz is one of the most abundant minerals and essential constituent of many igneous, sedimentary, and metamorphic rocks (Yao *et al.*, 2017; Del Lama and de Frascá, 2018). Quartz is a chemically stable mineral common in the soil environment but can be washed by runoff into the aquatic ecosystem (Baldermann *et al.*, 2021).

Studies revealed that identifying the source and constituents of SPM and

sediments is necessary for the establishment of the pollution status of aquatic system (Ma *et al.*, 2015, Zuliani *et al.*, 2022). Researches on the evaluation of minerals and metals in SPM and sediments have been carried out in countries outside Nigeria (Lučić *et al.*, 2018; Zhang *et al.*, 2018; Zeng *et al.*, 2020; Gordeev *et al.*, 2022). However, information on the geological and mineralogical of SPM and sediment in aquatic ecosystem in the study area are limited. Emphasis are more on the physicochemical, metals and microbial loads of water, SPM, and sediment samples from rivers within the region (Abu and Egenonu, 2008; Igbinsosa *et al.*, 2012; Ebong and Etuk, 2017; Ebong and John, 2021; Ehiemere *et al.*, 2022; Izah *et al.*, 2022; Iwegbue *et al.*, 2023).

The elemental and mineral compositions of SPM and sediments from some rivers within the Southeastern part of Nigeria were examined in this study. The objectives of this study were (i) to assess the source of metals in the studied SPM and sediments, (ii) to identify the minerals present in the studied samples, (iii) to understand the relationship between minerals constituents of sediment and concentrations of metals in the studied rivers, (iv) to provide a comprehensive information on the pollution status of the rivers investigated, and (v) to reveal the contributions of natural rock weathering processes and anthropogenic activities on the minerals and metals of these samples.

Materials and methods

Study area

The areas covered during this research lie between longitudes 7°31'E - 8°23'E and latitude 4°33'N - 5°10'N in the South-Eastern section of the Niger-Delta Region of Nigeria (Figure 1). Three (3) estuaries (Imo, Qua-Iboe, and Cross Rivers) and two (2) tidal rivers (Calabar and Great Kwa Rivers) were investigated in this study. The coordinates of the studied locations as obtained from the Geographical Positioning System (GPS) are in Table 1. The area investigated is a sedimentary basin consisting of Quaternary and Pleistocene sediments. The studied rivers are within the meso-tidal setting where the influence of tide and waves are experienced (Boboye and Akinmosin, 2018). The study area is within the subequatorial climatic zone in the humid tropical rainforest area (Hassan *et al.*, 2020). The studied area is characterized by two prominent seasons namely: Dry season which lasts from November to March and Wet season that starts from April to October (Adejuwon, 2012; Murtala *et al.*, 2020). The atmospheric temperature of the area is at its highest in February and lowest between July and September (Kemela and Phoebe, 2021). The humidity peaks in July while the minimum level is observed in the month of January (Ebong *et al.*, 2020).

Sample collection and treatment

Water and sediment samples were obtained from Great Kwa, Calabar, Cross River, Imo, and Qua Iboe Rivers

at high and low tides. The collection of water and sediment samples was done between November 2011 and February 2012. Water samples were obtained from the rivers investigated from the upper 20cm of the water channel in a 25-liter plastic containers free oil (Moussa *et al.*, 2022). Samples collected were homogenized and filtered through already dried and weighed Whatman fiberglass filters with a porosity of 0.45 mm. Sediments collected on the filter paper were dried again at 105°C for 90

minutes and re-weighed to achieve the desired SPM (Zhang *et al.*, 2014). The SPM obtained were stored in a cooler for further analyses. Bottom sediments were collected at the various rivers investigated at a depth of 0-5 cm using a Van Veen Grab sampler, transferred into nylon bags and transported in a cooler to the Laboratory. These sediment samples were air-dried at room temperature for seven (7) days (Rzetala *et al.*, 2019; Mackowiak and Perdrial, 2023).




Figure 1: Geological map of the study area showing sample locations.

Table 1: Site location and Coordinates.

S/ N	River System	Longitude	Latitude	Location
1	Great Kwa River	8° 23' 40.8516" E	4° 56' 57.9516" N	Apabuyo Bridge
2	Calabar River	8° 19' 21.9108" E	4° 58' 38.9208" N	Near Former CALCEMCO
3	Cross River	8° 4' 8.0112" E	5° 10' 43.1400" N	Calabar Itu Bridge
4	Imo River	7° 31' 15.8196" E	4° 40' 49.8216" N	At CrossRiver/Akwa Ibom Bridge
5	Qua Iboe River	7° 56' 46.2084" E	4° 33' 46.2384" N	Near Qua Iboe Town

Mineralogical analyses

The SPM and sediments obtained from the studied aquatic ecosystems were subjected to elemental and mineralogical compositions analyses as shown below. Quantitative compositional analysis of the samples was performed using a four spectrometer Cameca SX50 electron microprobe with a 10 nA beam current and accelerating voltage of 15 kV following the methods of Chebotarev *et al.* (2022) and Pérez-Huerta *et al.* (2023).

Quantitative analyses were performed with wavelength-dispersive spectrometers (WDS) after standardization with properly characterized compounds. Qualitative analyses (spectra) were obtained using an Imix Princeton Gamma Tech (PGT) energy dispersive system (EDS) using a thin-window detector (Stevenson *et al.*, 2000; Mackowiak and Perdrial, 2023). Mineral compositions of the samples were obtained using Energy Dispersion X-Ray Analysis (EDX) according to the methods of Han *et al.* (2022). Scanning electron microscope (SEM) uses a beam of high-energy electrons to generate signals at the surface of the solid specimen. Signals derived from the electron-sample interactions shows the orientation of the materials making up the sample (Han *et al.*, 2018; Lučić *et al.*, 2019).

All mineralogical measurements were done with X-ray diffraction (XRD) analysis on a Siemens D8 X-ray diffractometer with a Cu K α radiation over a 2 θ range of 5 to 60°, a step interval of 0.02°, and duration of 4s

step⁻¹ (Kravchishina and Dara, 2014; Ali *et al.*, 2022).

Results and discussion

Elemental and mineral compositions of suspended particulate matter and sediment from the studied rivers

Results of the compositions and levels of elements in sediments from the five (5) rivers investigated are indicated in Figures 2–6 below. Figures 7 and 8 illustrate the *mineralogical studies in SPM and sediments, respectively. Mineral compositions of sediment from Great Kwa, Calabar, Cross River, Imo, and Qua Iboe Rivers* are illustrated in Figures 9–13.

The elemental compositions of sediments from the studied rivers revealed the following: Samples from Great Kwa River showed high levels of Si, Al, Ca Mg and Fe with traces of Ti, Na and K (Fig. 2). Samples from Calabar River composed of high concentrations of Si, Al, K, and Fe with low levels of Mg (Fig. 3). Figure 4 indicates that sediments from Cross River contained elevated levels of Si, Al, Ca, K, Mg, and low concentration of Ti. High levels of Si, Al, and Fe with low levels of K and Ti were recorded for sediments from Imo River (Fig. 5). Qua Iboe River samples had elevated levels of Si, Al, Mg, and Fe but low concentrations of Ti and Ca (Fig. 6). The high concentrations of Si and Al reported in sediments from all the rivers investigated is consistent with the results obtained by Krivtsov *et al.* (2020).

EDX analysis revealed that majority of the particles identified in the studied

sediments were mainly silicate minerals (quartz, clays, micas, feldspars, and mafic minerals (Mg-Fe) silicates) (Kimeli *et al.*, 2021).

Results of the X-ray diffraction analysis (XRD) for the mineral

compositions in suspended particulate matters (SPM) and sediments are shown in Figures 7 and 8, respectively.




Figure 2: EDX spectrum of Great Kwa River sediment.




Figure 3: EDX spectrum of Calabar River sediment.




Figure 4: EDX spectrum of Cross River sediment.




Figure 5: EDX spectrum of Imo River sediment.




Figure 6: EDX spectrum of Qua Iboe River sediment.




Figure 7: The XRD pattern of the suspended particulate matter. Blue = Great Kwa River; Purple = Calabar River; Green = Cross River; Dark red = Imo; River; Black = Qua Iboe River.




Figure 8: The XRD pattern of the sediment samples. Blue=Great Kwa River sediment; Purple=Calabar River sediment; Green=Cross River sediment; Red=Imo River sediment; Black = Qua Iboe River sediment.




Figure 9: SEM image of Great Kwa River sediment. 1- Quartz; 2- Zircon ($ZrSiO_4$); 3- Amphibole/pyroxene; 4- Biotite (mica minerals); 5- k-feldspar; 6- Pyrite (FeS_2); 7- FeOH.

These Figures reveal that both SPM and the sediment samples were dominated by quartz (28.56%). This is similar to the results obtained by Lučić *et al.* (2019) and Jianet *al.* (2020) in their studies. Quartz was identified by the diffraction peaks at 4.21, 4.23, 3.33, 3.23, 2.44, and 1.81Å. As shown in Figure 7, the reflection peak at 3.33Å is more intense in samples from Great Kwa River than in

other sediments. Hornblende with relative abundance of 20.11% was identified by the reflection peaks at 9.94, 9.91, and 8.34Å. Hornblende was detected mainly in the samples from the Great Kwa, Cross River, and Calabar Rivers.




Figure 10: SEM image of Calabar River Sediment. 1- Quartz; 2 - Clay mineral (kaolinite, illite, FeO).




Figure 11: SEM image of Cross River Sediment. 1- Quartz; 2-Pyroxene; 3-White mica (muscovite); 4- K-Feldspar; 5-Mafic minerals.




Figure 12: SEM image of Imo River. 1- K-feldspar; 2- Clay mineral (kaolinite, illite, FeO); 3- FeS₂ (pyrite); 4-Na-feldspar.




Figure 13: SEM image of Qua Iboe River. 1- Quartz; 2- K-feldspar; 3- Clay mineral (kaolinite, illite, FeO); 4- Chlorite.

Gypsum (15.31%) was identified by the reflection peaks at 7.11, 2.88, 2.27, and 2.23Å. Gibbsite was identified primarily by reflection peaks at 4.79 and 4.81Å. Gibbsite constituted about 15.55% of the entire minerals found in samples from Qua Iboe and Great Kwa Rivers, similar to the results obtained by Sidibe and Yalcin (2019). Clay minerals (illite (2.7%), kaolinite (3.8%), and chlorite (3.8%) were identified by their characteristics basal X-ray diffraction peaks at 2.54, 4.44 and 4.97Å,

respectively. Calabar and Cross Rivers samples contained traces of calcite (3.3%) and was identified by the reflection peaks at 2.94Å. Potassium feldspar with relative abundance of 12.10% and was identified by reflection peaks at 3.69, 3.55, and 3.46Å.

The XRD results revealed that apart from Quartz, clays, micas, feldspars, and mafic minerals identified by EDX, other minerals such as gypsum, gibbsite, and hornblende were also components in the studied SPM and sediments. This is in

agreement with the results obtained in a study within Sava River in Croatia by Lučić *et al.* (2018). This result is consistent with the morphological characterization done on the samples with scanning electron microscope and energy dispersive X-ray analyzer. The XRD pattern obtained indicated that sediment samples were more crystalline than suspended particulate matter. The pattern also revealed that, samples obtained during high tide contained more quartz, gypsum and gibbsite minerals than the low tide samples. This is consistent with the report obtained in dust micro particles by Grytting *et al.* (2022). Quartz was the most common type of silica minerals in all the samples examined as indicated by the intense Si peak in EDX spectra. The grain surface characteristics (Conchoidal fractures and V-shaped pits) observed in the SEM micrographs (Figs. 2-6) distinguished quartz from other minerals identified (Madhavaraju *et al.*, 2021). Quartz is one of the most abundant minerals and it is an essential constituent of many igneous, sedimentary, and metamorphic rocks (Yao *et al.*, 2017; Del Lama and de Frascá, 2018). Quartz mineral emanates from the mechanical and chemical weathering of quartz rich rocks, and as a chemically stable mineral, it is common in the earth soil environment can easily be washed into the aquatic environment (Baldermann *et al.*, 2021). This may have resulted in the observed abundance in the suspended particulate matters and sediments from the studied Rivers.

The scanning electron microscope (SEM) images of sediments from Great

Kwa River in Figure 9 indicate quartz, zircon, amphibole, pyroxene, biotite (mica mineral), potassium feldspar, pyrite, and iron hydroxide as the major minerals. Calabar River sediments contained mainly quartz and clay minerals with many organic matters (Fig. 10). Quartz, pyroxene, white mica, potassium feldspar, and mafic minerals were the dominant minerals in sediments from Cross River (Fig. 11). Potassium feldspar, clay minerals, pyrite, and sodium feldspar were the principal minerals sediments from Imo River (Fig. 12). SEM images of sediments from Qua Iboe River in Figure 13 indicate quartz, potassium feldspar, clay minerals (kaolinite, illite, ferrous oxide (FeO) and chlorite as the major minerals. This is consistent with the report by Kumari and Mohan (2021) in a similar study.

Conclusion

The study revealed the mineral and elemental compositions of SPM and sediments from Great Kwa, Calabar, Cross River, Imo, and Qua Iboe Rivers, Nigeria. EDX analysis indicated the elements present in sediment from each river however; Al and Si were the common ones. It was observed that, most of the particles in the studied sediments were silicate minerals. XRD analysis on SPM and sediments from rivers investigated were dominated by quartz. The XRD results revealed that gypsum, gibbsite, and hornblende were also major components of the studied SPM and sediments. Scanning electron microscope (SEM) images of sediments from the different rivers studied showed

varied minerals for the different locations. However, the common minerals in sediments from the various rivers studied were quartz and clay minerals. The study has shown that apart from anthropogenic factor, the natural geological processes in the aquatic environment can also contribute significant amounts of contaminants/pollutants to the studied rivers. The results obtained revealed the mineral and elemental compositions of the studied SPM and sediments.

References

- Abu, G.O. and Egenonu, C., 2008.** The current pollution status of the new Calabar River in the Niger Delta region of Southern Nigeria: A survey of antibiogram profiles of its bacterial isolates. *African Journal of Environmental Science and Technology*, 2(6), 134-141.
- Adejuwon, J.O., 2012.** Rainfall seasonality in the Niger Delta Belt, Nigeria. *Journal of Geography and Regional Planning*, 5(2), 51-60. DOI:10.5897/JGRP11.096
- Akhtar, N., Syakir Ishak, M.I., Bhawani, S.A. and Umar, K., 2021.** Various natural and anthropogenic factors responsible for water quality degradation: A review. *Water*, 13(19), 2660. DOI:10.3390/w13192660
- Algül, F. and Beyhan, M., 2020.** Concentrations and sources of heavy metals in shallow sediments in Lake Bafa, Turkey. *Scientific Reports*, 10, 11782. DOI:10.1038/s41598-020-68833-2
- Ali, A., Chiang, Y.W. and Santos, R.M., 2022.** X-ray Diffraction techniques for mineral characterization: A review for engineers of the fundamentals, applications, and research directions. *Minerals*, 12, 205. DOI:10.3390/min12020205.
- Baba, A. and Gündüz, O. 2017.** Effect of geogenic factors on water quality and its relation to human health around Mount Ida, Turkey. *Water*, 9(1), 66. DOI:10.3390/w9010066
- Baldermann, A., Dietzel, M. and Reinprecht, V., 2021.** Chemical weathering and progressing alteration as possible controlling factors for creeping landslides. *Science of the Total Environment*, 778, 146300. DOI:10.1016/j.scitotenv.2021.146300.
- Bartoszek, L., Gruca-Rokosz, R., Pękala, A. and Czarnota, J., 2022.** Heavy metal accumulation in sediments of small retention reservoirs. *Ecological risk and the impact of humic substances distribution. Resources*, 11,113. DOI:10.3390/resources11120113
- Bebetidoh, O.L, Kometa, S., Pazouki, K. and Norman, R., 2020.** Sustained impact of the activities of local crude oil refiners on their host communities in Nigeria. *Heliyon*, 5: 6(6), e04000. DOI:10.1016/j.heliyon.2020.e04000.
- Biscaye, P.E., Grousset, F.E., Revel, M., Van der Gaast, S., Zielinski, G.A., Vaars, A. and Kukla, G., 1997.** Asian provenance of glacial dust (stage 2) in the Greenland Ice Sheet Project 2 Ice Core, Summit, Greenland. *Journal of Geophysical Research*, 102(C12), 26765 – 26781.
- Boboye, O. and Akinmosin, A., 2018.** Palaeoclimatic evidences from the quaternary coastal deposits, southwestern, Nigeria. *Open Journal of Geology*, 8, 585-605. DOI:10.4236/ojg.2018.86034.
- Chebotarev, D.A., Wohlgenuth-Ueberwasser, C. and Hou, T., 2022.**

- Partitioning of REE between calcite and carbonatitic melt containing P, S, Si at 650–900 °C and 100 MPa. *Scientific Reports*, 12, 3320. DOI: 10.1038/s41598-022-07330-0
- Chen, J., Bouchez, J., Gaillardet, J. and Louvat, P., 2014.** Behaviors of major and trace elements during single flood event in the Seine River, France. *Procedia Earth and Planetary Science*, 10, 343-348.
- Chmielowska, D. and Salata, D., 2020.** Heavy minerals as indicators of the source and stratigraphic position of the loess-like deposits in the Orava Basin (Polish Western Carpathians). *Minerals*, 10, 445. DOI:10.3390/min10050445
- Del Lama, E.A. and de Frascá, M.H.B.O., 2018.** Metamorphic rocks. In: Bobrowsky, P.T., Marker, B. (eds) *Encyclopedia of Engineering Geology. Encyclopedia of Earth Sciences Series*. Springer, Cham. DOI:10.1007/978-3-319-73568-9_198
- Ebong, G.A., Ita, B.N., Nyong, A.B. and NU Benson, N.U., 2006.** Seasonal changes in the water quality of Qua Iboe River Estuary and its associated creeks in Ibeno, Nigeria. *Journal of Applied Sciences*, 9 (2), 6469-6482.
- Ebong, G.A. and Etuk, H.S. 2016.** Fractionation status and uptake potentials of trace metals in water bodies within Niger Delta Region, Nigeria. *International Journal of Pure & Applied Chemistry*, 13(2), 1-18. DOI:10.9734/IRJPAC/2016/29408
- Ebong, G.A. and Etuk, H.S., 2017.** Impact of anthropogenic activities on metal load of Sagbama River, Niger Delta Region of Nigeria. *Chemistry Research Journal*, 2(2), 96-103.
- Ebong, G.A. and John, R.C., 2021.** Physicochemical properties, total hydrocarbon content, and trace metals of water and sediments from major River Estuaries within the Niger Delta Region of Nigeria. *World Journal of Advanced Research and Reviews*, 12(02), 587–597. DOI:10.30574/wjarr.2021.12.2.0650
- Ebong, G.A., Ettesam, E.S. and Dan, E.U., 2020.** Impact of abattoir wastes on trace metal accumulation, speciation, and human health-related problems in soils within southern Nigeria. *Air, Soil and Water Research*, 13, 1–14. DOI:10.1177/117862211989843
- Ehiemere, V.C., Ihedioha, J.N., Ekere, N.R., Ibeto, C.N. and Abugu, H.O., 2022.** Pollution and risk assessment of heavy metals in water, sediment and fish (*Clarias gariepinus*) in a fish farm cluster in Niger Delta region, Nigeria. *J Water Health*, 20(6), 927–945. DOI:10.2166/wh.2022.003
- Ewim, D.R.E., Orikpete, O.F., Scott, T.O., Onyebuchi, C.N., Onukogu, A.O., Uzougbo, C.G. and Onunka, C., 2023.** Survey of wastewater issues due to oil spills and pollution in the Niger Delta area of Nigeria: a secondary data analysis. *Bulletin of the National Research Centre*, 47, 116. DOI:10.1186/s42269-023-01090-1
- Fan, J., Jian, X., Shang, F., Zhang, W., Zhang, S. and Fu, H., 2021.** Underestimated heavy metal pollution of the Minjiang River, SE China: Evidence from spatial and seasonal monitoring of suspended-load sediments. *Science of the Total Environment*, 760, 142586. DOI:10.1016/j.scitotenv.2020.142586
- Fan, J., Yang, J., Cheng, F. and Zhang, S., 2023.** The source, distribution, and environmental effects of suspended particulate matter in the Yangtze River

- System. *Water*, 15, 3429. DOI: 10.3390/w15193429
- Fernandes, L.L., Kessarkar, P.M., V. Roa, V.P., Suja, S., Parthiban, G. and Kurian, S., 2019.** Seasonal distribution of trace metals in suspended particulate and bottom sediments of four microtidal river estuaries, west coast of India. *Hydrological Sciences Journal*, 64,12, 1519-1534. DOI:10.1080/02626667.2019.1655147
- Gordeev, V.V., Dara, O.M., Filippov, A.S., Belorukov, S.K., Lokhov, A.S., Kotova, E.I. and Kochenkova, A.I., 2022.** Mineralogy of particulate suspended matter of the severnaya Dvina River (White Sea, Russia). *Minerals*, 12, 1600. DOI:10.3390/min12121600
- Gordon, B.A., Dorothy, O. and Lenhart, C.F., 2020.** Nutrient retention in ecologically functional floodplains: a review. *Water (Basel)*, 12, 2762. DOI:10.3390/w12102762
- Grytting, V.S., Refsnes, M., Låg, M., Erichsen, E., Røhr, T.S., Snilsberg, B., White, R.A. and Øvrevik, J., 2022.** The importance of mineralogical composition for the cytotoxic and pro-inflammatory effects of mineral dust. *Particle and Fibre Toxicology*, 19, 46. DOI:10.1186/s12989-022-00486-7
- Han, W., Jiao, H. and Fox, D., 2018.** *Scanning Electron Microscopy*. In: Wang, R., Wang, C., Zhang, H., Tao, J., Bai, X. (eds) Progress in nanoscale characterization and manipulation. Springer Tracts in Modern Physics, vol 272. Springer, Singapore. DOI:10.1007/978-981-13-0454-5_2
- Han, S., Löhr, S.C., Abbott, A.N., Baldermann, A., Farkaš, J., McMahon, W., Milliken, K.L., Rafiei, M., Wheeler, C. and Owen, M., 2022.** Earth system science applications of next-generation SEMEDS automated mineral mapping. *Frontiers of Earth Science*, 10, 956912. DOI:10.3389/feart.2022.956912
- Hassan, I., Kalin, R. M., Aladejana, J. and White, C.J., 2020.** Potential impacts of climate change on extreme weather events in the Niger Delta Part of Nigeria. *Hydrology*, 7(1),1-19. DOI:10.3390/hydrology7010019
- Hope, J.A., Hewitt, J., Pilditch, C.A., Savage, C. and Thrush, S.F., 2020.** Effect of nutrient enrichment and turbidity on interactions between microphytobenthos and a key bivalve: Implications for higher trophic levels. *Frontiers in Marine Science*, 7, 695. DOI:10.3389/fmars.2020.00695
- Igbinsosa, E.O., Uyi, O.O., Odjadjare, E.E., Ajuzie, C.U., Orhue, P.O. and Adewole, E.M., 2012.** Assessment of physicochemical qualities, heavy metal concentrations and bacterial pathogens in Shanomi Creek in the Niger Delta, Nigeria. *African Journal of Environmental Science and Technology*, 6(11), 419-424. DOI:10.5897/AJEST12.038
- Iwegbue, C.M.A., Faran, T.K., Iniaghe, P.O., Ikefan, J.O., Tesi, G.O., Nwejei, G.E. and Martincigh, B.S., 2023.** Water quality of Bomadi Creek in the Niger Delta of Nigeria: Assessment of some physicochemical properties, metal concentrations, and water quality index. *Applied Water Science*, 13, 36. DOI:10.1007/s13201-022-01804-2
- Izah, S.C., Ayobami O. Aigberua, A.O. and Arun Lal Srivastav, A.L., 2022.** Chapter 4 - Factors influencing the alteration of microbial and heavy metal characteristics of river systems in the Niger Delta region of Nigeria. Editor(s), Madhav, S., Kanhaiya, S., Srivastav, A.,

- Singh, V. and Singh, P. Ecological Significance of River Ecosystems, Elsevier, pp. 51-78. DOI:10.1016/B978-0-323-85045-2.00005-4
- Jian, X., Zhang, W., Yang, S. and Kao, S.J., 2020.** Climate-dependent sediment composition and transport of mountainous rivers in tectonically stable, subtropical east Asia. *Geophysical Research Letters*, 47(3), e2019GL086150. DOI:10.1029/2019GL086150
- Kemela, B. and Phoebe, A., 2021.** Assessment of climate variability in the Niger Delta Region of Nigeria. *International Journal of Chemistry and Chemical Engineering Systems*, 6, 17 – 28. <http://www.ias.org/ias/journals/ijces>
- Kimeli, A., Ocholla, O., Okello, J., Koedam, N., Westphal, H. and Kairo, J., 2021.** Geochemical and petrographic characteristics of sediments along the transboundary (Kenya–Tanzania) Umba River as indicators of provenance and weathering. *Open Geosciences*, 13(1),1064-1083. DOI:10.1515/geo-2020-0275
- Kloprogge, J.T. and Wood, B.J., 2020.** Handbook of Mineral Spectroscopy Volume 1. X-Ray Photoelectron Spectra; Elsevier: Amsterdam, The Netherlands, 505P.
- Kravchishina, M.D. and Dara, O.M., 2014.** Mineral composition of the suspended particulate matter in the White Sea. *Oceanology*, 54(3), 327–337.
- Krivtsov, V., Arthur, S., Buckman, J., Kraiphet, A., Needham, T., Gu, W., Gogoi, P. and Thorne, C., 2020.** Characterization of suspended and sedimented particulate matter in blue-green infrastructure ponds. *Blue-Green Systems*, 2(1), 214–236. DOI:10.2166/bgs.2020.102
- Kumari, N. and Mohan, C., 2021.** Basics of clay minerals and their characteristic properties. *IntechOpen*. DOI:10.5772/intechopen.97672
- Lednicka, B., Kubacka, M., Freda, W., Haule, K., Dembska, G., Galer-Tatarowicz, K. and Pazikowska-Sapota, G., 2022.** Water Turbidity and Suspended Particulate Matter Concentration at Dredged Material Dumping Sites in the Southern Baltic. *Sensors*, 22(20),8049. DOI:10.3390/s22208049
- Li, Y., Wang, X., Fu, W., Xia, X., Liu, C., Min, J., Zhang, W. and Crittenden, J.C., 2019.** Interactions between nano/micro plastics and suspended sediment in water: implications on aggregation and settling. *Water Research*, 161, 486-495. DOI:10.1016/j.watres.2019.06.018
- Lin, L., Yang, H. and Xu, X., 2022.** Effects of Water Pollution on Human Health and Disease Heterogeneity: A Review. *Frontiers of Environmental Science*, 10:880246. DOI:10.3389/fenvs.2022.880246
- Lučić, M., Jurina, I., Ščančar, J., Mikac, N. and Vdović, N., 2018.** Sedimentological and geochemical characterization of river suspended particulate matter (SPM) sampled by time-integrated mass flux sampler (TIMS) in the Sava River (Croatia). *Journal of Soils and Sediments*. DOI:10.1007/s11368-018-2104-2

- Lučić, M., Jurina, I., Ščančar, J., Mikac, N. and Vdović, N., 2019.** Sedimentological and geochemical characterization of river suspended particulate matter (SPM) sampled by time-integrated mass flux sampler (TIMS) in the Sava River (Croatia). *Journal of Soils and Sediments*, 19, 989–1004. DOI:10.1007/s11368-018-2104-2
- Luvhimbi, N., Tshitangano, T.G., Mabunda, J.T., Olaniyi, F.C. and Edokpayi, J.N., 2022.** Water quality assessment and evaluation of human health risk of drinking water from source to point of use at Thulamela municipality, Limpopo Province. *Scientific Reports*, 11, 12(1),6059. DOI:10.1038/s41598-022-10092-4.
- Ma, Y., Qin, Y., Zheng, B., Zhang, L. and Zhao, Y., 2015.** Seasonal variation of enrichment, accumulation and sources of heavy metals in suspended particulate matter and surface sediments in the Daliao river and Daliao river estuary, Northeast China. *Environmental Earth Sciences*, 73, 5107-5117.
- Macheyeki, A.S., Li, X., Kafumu, D.P. and Yuan, F., 2020.** Chapter 1 - *Elements of exploration geochemistry*. Applied Geochemistry, Elsevier, pp. 1-43. DOI:10.1016/B978-0-12-819495-9.00001-3
- Mackowiak, T.J. and Perdrial, N., 2023.** Monitoring of Suspended Sediment Mineralogy in Puerto Rican Rivers: Effects of Flowrate and Lithology. *Minerals*, 13(2), 208. DOI:10.3390/min13020208
- Madhavaraju, J., John S. Armstrong-Altrin, J.S., James, R.A. and Hussain, S.M., 2021.** Palaeoenvironment and provenance signatures inferred from quartz grain surface features: A case study from Huatabampo and Altata beaches, Gulf of California, Mexico. *Journal of South American Earth Sciences*, 111, 103441. DOI:10.1016/j.jsames.2021.103441
- Magana-Arachchi, D.N. and Wanigatunge, R.P., 2020.** Ubiquitous waterborne pathogens. *Waterborne Pathogens*, 15–42. DOI:10.1016/B978-0-12-818783-8.00002-5
- Moussa, M.B., Touré, A.A., Kergoat, L., Lartiges, B., Rochelle-Newall, E., Robert, E., Gosset, M., Tanimoun, B.A. and Grippa, M., 2022.** Spatio-temporal dynamics of suspended particulate matter in the middle Niger River using in-situ and satellite radiometric measurements. *Journal of Hydrology: Regional Studies*, 41, 101106. DOI:10.1016/j.ejrh.2022.101106
- Murtala, A.I., Oguntunde, P.G., Akinwumiju, A.S. and Olubanjo, O.O., 2020.** Rainfall analysis over the Niger Central Hydrological Area, Nigeria: Variability, Trend, and Change point detection. *Scientific African*, 8, e00419. DOI:10.1016/j.sciaf.2020.e00419
- Nair, N.V. and Nayak, P.K., 2023.** Exploring water quality as a determinant of small-scale fisheries vulnerability. *Sustainability*, 15, 13238. DOI: 10.3390/su151713238
- Ouyang, N., Zhang, Y., Sheng, H., Zhou, Q., Huang, Y. and Yu, Z., 2021.** Clay mineral composition of upland soils and its implication for pedogenesis and soil taxonomy in subtropical China. *Sci Rep.*, 11:9707. DOI: 10.1038/s41598-021-89049-y
- Patrolecco, L., Ademollo, N., Capri, S., Pagnotta, R. and Polesello, S., 2010.** Occurrence of priority hazardous PAHs in water, suspended particulate matter,

- sediment and common eels (*Anguilla anguilla*) in the urban stretch of the River Tiber (Italy). *Chemosphere*, 81, 1386-1392.
10.1016/j.chemosphere.2010.09.027
- Pérez-Huerta, A., Brugal, J.P., Salomé, M., Schmitt, C.N.Z. and Dauphin, Y., 2023.** Integrated information on the structure and composition of the ostrich eggshell (*Struthio camelus*). *Minerals*, 13, 481. DOI: 10.3390/min13040481
- Pye, K. and Blott, S.J., 2006.** Coastal processes and morphological change in the Dunwich - Sizewell area, Suffolk, UK. *Journal of Coastal Research*, 22, 453-473. <http://113.160.249.209:8080/xmlui/handle/123456789/2968>
- Rzetala, M., Babicheva, V.A. and Rzetala, M.A., 2019.** Composition and physico-chemical properties of bottom sediments in the southern part of the Bratsk Reservoir (Russia). *Scientific Reports*, 9(1), 12790.
- Saito, K., Tada, R., Zheng, H., Irino, T., Luo, C., He, M., Wang, K. and Suzuki, Y., 2017.** ESR signal intensity of quartz in the fine-silt fraction of riverbed sediments from the Yangtze River: a provenance tracer for suspended particulate matter. *Progress in Earth and Planetary Sciences*, 4:4. DOI: 10.1186/s40645-017-0118-9
- Shil, S. and Singh, U.K., 2019.** Health risk assessment and spatial variations of dissolved heavy metals and metalloids in a tropical river basin system. *Ecological Indicators*, 106, 105455. DOI: 10.1016/j.ecolind.2019.105455.
- Sidibe, M. and Yalcin, M.G., 2019.** Petrography, mineralogy, geochemistry and genesis of the Balaya bauxite deposits in Kindia region, Maritime Guinea, West Africa. *Journal of African Earth Sciences*, 149, 348-366. DOI: 10.1016/j.jafrearsci.2018.08.017.
- Sojka, M. and Jaskula, J., 2022.** Heavy Metals in River Sediments: Contamination, toxicity, and source identification-A case study from Poland. *International Journal of Environmental Research and Public Health*, 19(17), 10502. DOI: 10.3390/ijerph191710502
- Stevenson, D.S., Johnson, C.E., Collins, W.J., Derwent, R.G. and Edwards, J.M., 2000.** Future estimates of tropospheric ozone radiative forcing and methane turnover — The impact of climate change. *Geophys. Res. Lett.*, 27, 2073–2076. DOI: 10.1029/1999GL010887
- Sun, G., Wang, M., Guo, J., Wang, Y. and Yang, Y., 2020.** Geochemical significance of clay minerals and elements in paleogene sandstones in the center of the northern margin of the Qaidam Basin, China. *Minerals*, 10(6), 505. DOI: 10.3390/min10060505
- Surette, M.C. and Nason, J.A., 2019.** Nanoparticle aggregation in a freshwater river: the role of engineered surface coatings. *Environmental Science: Nano*, 6, 540-553. DOI:10.1039/C8EN01021H
- Sutherland, T.F., Amos, C.L. and Grant, J., 2022.** The resuspension and deposition of biomediated sediments in upper south Cove, Nova Scotia, Canada. *Journal of Coastal Research*, 38(1), 19-34. <https://www.jstor.org/stable/48646239>
- Tian, S., Li, Z., Wang, Z., Jiang, E., Wang, W. and Sun, M., 2021.** Mineral composition and particle size distribution of river sediment and loess in the middle and lower Yellow River. *International*

- Journal of Sediment Research*, 36(3),392-400. DOI: 10.1016/j.ijsrc.2020.07.008.
- Vidmar, J., Zuliani, T., Novak, P., Drinčić, A., Ščančar, J. and Milačić, R., 2017.** Elements in water, suspended particulate matter and sediments of the Sava River. *Journal of Soils and Sediments*, 17, 1917-1927. DOI:10.1007/s11368-016-1512-4
- Wagner, S., Hüffer, T., Klöckner, P., Wehrhahn, M., Hofmann, T. and Reemtsma, T., 2018.** Tire wear particles in the aquatic environment - A review on generation, analysis, occurrence, fate and effects. *Water Research*, 139, 83-100. DOI:10.1016/j.watres.2018.03.051
- Walch, H., von der Kammer, F. and Hofmann, T., 2022.** Freshwater suspended particulate matter—Key components and processes in floc formation and dynamics. *Water Research*, 220, 118655. DOI: 10.1016/j.watres.2022.118655.
- Yao, K., Pradhan, B. and Idrees, M.O., 2017.** Identification of rocks and their quartz content in gua musang goldfield using advanced space borne thermal emission and reflection radiometer imagery. *Journal of Sensors*,6794095. DOI: 10.1155/2017/6794095
- Yin, S., Yi, Y., Liu, Q., Luo, Q. and Chen, K., 2022.**A review on effects of human activities on aquatic organisms in the Yangtze River Basin since the 1950s. *River*, 1(1),104-119. DOI: 10.1002/rvr2.15
- Yoshimura, T., Araoka, D., Kawahata, H., Hossain, H.M.Z. and Ohkouchi, N., 2021.** The influence of weathering, water Sources, and hydrological cycles on lithium isotopic compositions in river water and groundwater of the Ganges–Brahmaputra–Meghna River system in Bangladesh. *Frontiers of Earth Science*, 9, 668757. DOI:10.3389/feart.2021.668757
- Zeng, J., Han, G., Wu, Q. and Tang, Y., 2019.** Heavy metals in suspended particulate matter of the Zhujiang River, southwest China: Contents, sources, and health risks. *International Journal of Environmental Research and Public Health*, 16(10), 1843. DOI: 10.3390/ijerph16101843
- Zeng, J., Han, G. and Yang, K., 2020.** Assessment and sources of heavy metals in suspended particulate matter in a tropical catchment, northeast Thailand. *Journal of Cleaner Production*, 265: 121898. DOI: 10.1016/j.jclepro.2020.121898
- Zhang, J., Zhou, F., Chen, C., Sun, X., Shi, Y., Zhao, H. and Chen, F., 2018.** Spatial distribution and correlation characteristics of heavy metals in the seawater, suspended particulate matter and sediments in Zhanjiang Bay, China. *PLoS ONE*, 13(8), e0201414. DOI: 10.1371/journal.pone.0201414
- Zhang, M.W., Dong, Q., Cui, T.W., Xue, C.J. and Zhang, S.L., 2014.** Suspended sediment monitoring and assessment 587 for Yellow River estuary from Landsat TM and ETM plus imagery. *Remote Sensing of Environment*, 146, 588 136-147. DOI: 10.1016/j.rse.2013.09.033
- Zuliani, T., Vidmar, J., Ščančar, J., Kolarević, M.K., Kolarević, S., Paunović, M. and Milačić, R., 2022.** Transport of Potentially Toxic Elements in Solid Particulate Matter during Flash Flood Events in Upper and Lower Stretch of the Sava River. *Water*, 14(8),1213. DOI: 10.3390/w14081213