



## A Historical Analysis of Aquatic Research Threats

**Naim Oblomurodov<sup>1\*</sup>; Abdumajid Madraimov<sup>2</sup>; Zulfiya Palibayeva<sup>3</sup>;  
Askariy Madraimov<sup>4</sup>; Muhammadjon Zufarov<sup>5</sup>; Mushtariy Abdullaeva<sup>6</sup>;  
Bakhtiyor Pardaev<sup>7</sup>; Kurbonaliyon Zokirov<sup>8</sup>**

Received: 05 August 2024; Revised: 05 September 2024; Accepted: 04 November 2024; Published: 10 December 2024

### Abstract

In the field of marine sciences, historical study is becoming more and more significant. The problem of moving baselines for some species and biological systems has been tended to by the utilization of verifiable information in marine asset the executives and policymaking. Huge headways in calculated and systemic techniques are expected to add to an exhaustive comprehension of the worldwide history of human cooperations with marine life, even if many significant research problems remain unsolved. This study looks at how dangers to aquatic research have changed over time, starting in the 19th century. A complex web of interrelated causes has weakened aquatic ecosystems' health and biodiversity, according to the report. Aquatic ecosystems' health and biodiversity have been jeopardized by environmental degradation, overfishing, a lack of financing, multidisciplinary conflicts, and regulatory difficulties. This analysis emphasizes the necessity of a multidisciplinary strategy to combat these risks as well as ongoing funding for conservation and research. This analysis emphasizes the necessity of a multidisciplinary strategy to tackle the challenges confronting aquatic research and the significance of sustained investment in conservation and research initiatives to safeguard aquatic ecosystems worldwide.

**Keywords:** Historical research, Marine science, Aquatic research, Environmental degradation.

---

1\*- Tashkent State University of Economics, Uzbekistan. Email: oblomurodov@gmail.com,

ORCID: <https://orcid.org/0009-0002-5353-7218>

2- State Museum of the History of Temurids of the Academy of Sciences of Uzbekistan, Uzbekistan.

Email: abdulmajidmadraimov@gmail.com, ORCID: <https://orcid.org/0009-0001-2969-0282>

3- State Conservatory of Uzbekistan Institute for Training Specialists in Paralympic Sports, Tashkent,

Uzbekistan. Email: p.zulfiya@mail.ru, ORCID: <https://orcid.org/0009-0003-0933-6301>

4- Tashkent State University of Oriental Studies, Uzbekistan. Email: askariy\_madraimov@tsuos.uz,

ORCID: <https://orcid.org/0009-0000-2238-2888>

5- Tashkent State Law University, Uzbekistan. Email: muhammadjonzufarov2@gmail.com,

ORCID: <https://orcid.org/0009-0009-4234-4008>

6- National University of Uzbekistan named Mirzo Ulugbek Tashkent, Uzbekistan.

Email: abdullaeva\_mushtariy@nuu.uz, ORCID: <https://orcid.org/0009-0007-6943-3663>

7- Jizzakh State Pedagogical University, Uzbekistan. Email: baxtiyorpardayev89@gmail.com,

ORCID: <https://orcid.org/0000-0001-6868-753X>

8- Tashkent State Agrarian University, Uzbekistan. Email: k\_zokirov@tdau.uz,

ORCID: <https://orcid.org/0000-0002-8156-5913>


\*Corresponding author

DOI: 10.70102/IJARES/V4S1/2

## Introduction

It is believed that the Phoenicians, who came from an ancient Mediterranean empire, may have been the first to study marine biology as early as 1200 BC. Aristotle was the first to document observations of aquatic life. Captain James Cook's exploration marked the beginning of the modern study of marine biology (SV and Bhandarkar, 2013). Raising global familiarity with the significance of submerged social legacy and including leaders during the time spent safeguarding these social fortunes are basic given the flood in interest in undersea social legacy throughout the course of recent many years. Underwater cultural heritage sites may be threatened by environmental changes like climate change, increased erosion, and current shifts (Tiwari *et al.*, 2017). However, we can learn a lot about historical climate change that formerly affected our ancestors' lives via undersea cultural legacy (Vollenweider *et al.*, 2003). Sea levels have been rising because of environmental change; this is a logical truth. Ocean level variances on the mainland edge are brought about by the collaboration between rising ocean levels and the upward uprooting of the world's outside layer. Planning and carrying out archaeological surveys on the continental shelf therefore requires taking these changes into consideration. Because of altered conservation patterns, shifting flows, and the presentation of new creature species into waters, climate change can potentially result in the destruction of several sites (Saha and Zaman, 2011). Several basic features of aquatic ecosystems may help to explain

the scientific breakthroughs that have resulted from extensive environmental research on these systems. First, compared to the earthbound climate, significant cycles in sea-going environments happen all the more rapidly. In contrast to yearly or lasting plants in earthbound frameworks, the essential and optional makers, including green growth and invertebrate slow eaters, develop and kick the bucket quickly (frequently between hours to months). Second, since amphibian environments are found at low rises in the scene and accumulate material from the watershed and air shed, they can offer integrative appraisals of territorial movement. Third, dissimilar to earthbound biology, which has generally isolated creature and plant environment, sea-going examinations have generally been integrative information on abiotic variables, creatures, and essential makers are often accumulated all the while (Sullieva *et al.*, 2024). Fourth, lakes enjoy a one of a kind upper hand over other sea-going environments: their effectively noticeable boundaries make a moderately restricted framework that takes into consideration the investigation of the multitude of pinions and wheels of energy stream. Lastly, human cultures depend on marine and freshwater systems for easily accessible water, wholesome food, transportation routes, leisure activities, and cultural identity components (Mateo-Sagasta *et al.*, 2018). A diagram summarizing threats to aquatic biodiversity and corresponding management strategies shown in Figure 1.



**Figure 1: A diagram summarizing threats to aquatic biodiversity and corresponding management strategies.**

### General Threats of Aquatic Ecosystem

The need of checking these regular assets for the administration and protection of these environment administrations is the premise of some drawn out oceanic investigations. Past these inherent properties of environments is a common trait of practically throughout the entire fruitful term studies — persistent people who figure out how to make long haul financing with momentary awards, energetic by the logical secrets unfurling, the cultural significance of the main things in need of attention, and habitually by the grandness of the biological system itself. Oceanic biological systems are settings that depend on water and where biotic and abiotic components communicate (Appenroth, 2010). Regularly, "oceanic environments" are isolated into two classifications:

freshwater and marine. With over 70% of the World's surface covered by water, the marine biological system is the biggest. Seas, estuaries, coral reefs, and waterfront biological systems are the few classifications that make up the marine environment. Under 1% of the World's surface is comprised of freshwater conditions. Lotic, lentic, and wetland environments are the different types of freshwater ecosystems. When an aquatic ecosystem's capacity to withstand stress is exceeded, its health deteriorates. Environmental changes that are physical, chemical, or biological can cause stress on an aquatic ecosystem. Changes in water temperature, velocity, and light availability are examples of physical changes. Changes in the loading rates of poisons, oxygen-consuming substances, and bio stimulatory nutrients are

examples of chemical changes. Exotic species introduction and excessive commercial species harvesting are examples of biological changes. Aquatic habitats may experience severe stress due to human populations.

### **Human Exercises Bringing about Tainting of Sea-going Biological Systems**

Aquatic habitats become contaminated as a result of anthropogenic exercises such as deforestation, filling and building waterways, dams, streets, and scaffolds, as well as farming, modern, and residential operations. The primary causes of water contamination are agriculture, industry, and human settlements.

**Agrochemicals:** The growing demand for food has resulted in the expansion of agriculture and the clearing of land, both of which have increased the levels of pollution in the water. Growing populations have raised the need for food, which has led to a rise in the amount of agrochemicals used to boost output (Lamberth *et al.*, 2013). The ecosystem, counting streams, lakes, springs, and beach front waters, is turning out to be more dirtied because of the unreasonable utilization of agrochemicals (manures, pesticides, herbicides, and plant chemicals) to increment creation.

**Nutrients:** Water pollution occurs when fertilizers are administered more frequently than the soil can fix them, absorb them by the crops, or remove them through surface overflow from the dirt. Abuse of phosphate and nitrogenous composts can enter surface water bodies through surface overflow or break into groundwater.

**Salts:** As a result of irrigation, deposited salts in the soil are carried by drainage water into receiving water bodies, where they lead to salinization. Another significant factor contributing to salinization in coastal regions is the infiltration of salty seawater into groundwater springs because of unnecessary groundwater extraction for horticultural purposes. The geochemical patterns of significant components like carbon, iron, nitrogen, phosphorus, silicon, and sulfur are altered by highly salinized waters, which has an overall effect on ecosystems. By altering species and community composition and lowering the richness of microbes, algae, plants, and animals, salinization can have an impact on freshwater biota.

**Emerging Pollutants:** Over the past 20 years, new agricultural pollutants have appeared, including hormones, growth boosters, vaccinations, and antibiotics. These contaminants can enter waterways by slurries and manure applied to agricultural land, as well as through runoff and leaching from aquaculture and animal operations. Currently, aquatic habitats in Europe are known to include around 700 new contaminants, together with their metabolites and transformation products. By using wastewater for irrigation and applying municipal biosolids to land as fertilizer, agriculture not only creates new pollutants but also aids in their spread and reintroduction into aquatic habitats. Attention must be paid to the possible health dangers to people caused by exposure to newly discovered toxins through tainted agricultural goods.

**Sewage:** Sewage is the largest type of waste that is released into aquatic environments. Industrial, municipal, and

household wastes, such as those from kitchens, bathrooms, washing machines, and feces, are all found in sewage. The optimum sinks for the discharge of these pollutants are freshwater sources. Clean water resources are being depleted and water contamination has increased as a result of sewage releases. The waste frameworks, which commonly open into neighboring waterways or sea-going frameworks, in the long run wash off the huge amounts of such junk that are delivered consistently from thickly populated urban communities. Various natural issues have been brought about by it, remembering a decrease for the sum and nature of water, serious flooding, eradication of species, and changes to the circulation and construction of the oceanic (Bhat and Qayoom, 2021).

### Uzbekistan Aquatic Threats

**Water scarcity:** More than 80% of Uzbekistan's water comes from its neighbors, making it reliant on them. By 2030, the nation's water deficit might be 7 billion cubic meters, and by 2050, it might be 14 billion. **Degradation of the environment:** The Aral Sea has vanished as a result of excessive irrigation and river flow allocation. The expanded utilization of pesticides in farming has additionally changed the nature of groundwater and surface water. **Decline of native fish species:** because of environmental change and human movement, Uzbekistan has seen a sharp reduction in the number of endemic fish species. Some species are probably extinct now, such as the Samarkand char.

- **Qosh Tepa Canal:** The Taliban's development of this canal may jeopardize Turkmenistan's and

Uzbekistan's water security. Up to one-third of the water in the Amu Darya could be redirected by the canal.

- **Old Irrigation System:** It is believed that 30–60% of water is lost due to Uzbekistan's outdated irrigation infrastructure. Human activity is primarily to blame for the deterioration of aquatic environments.
- Degradation of the environment includes pollution, ocean acidification, habitat loss, and climate change.
- Bycatch, discarding, and unregulated fishing practices are examples of overfishing and exploitation.
- **Lack of Finance and Resources:** insufficient funding, limited access to research facilities, and brain drain
- **Interdisciplinary Conflicts:** Disagreements among stakeholders, policymakers, and scientific fields.
- **Issues with Regulations and Policies:** insufficient rules, non-enforcement, and contradictory policies.
- **Ocean Acidification and Climate Change:** escalating temperatures, increasing sea levels, and modifications to ocean chemistry
- **Invasive Species and Diseases:** Introduction of non-native species and diseases that harm aquatic ecosystems.

Water contamination is mostly caused by increased industrialization and urbanization. The disposal of solid waste,

industrial waste, and household garbage are only a few of the many ways that humans contribute to water contamination. The globe is very concerned about water pollution. To lessen the contamination of aquatic habitats, environmental education is crucial.

### Conclusion

The health and biodiversity of aquatic ecosystems have been jeopardized by a complex and linked web of variables, as this historical review of aquatic research challenges demonstrates. These risks have changed throughout time and still affect aquatic research and conservation initiatives, ranging from overfishing and environmental degradation to financial constraints and regulatory obstacles. The study's conclusions emphasize the necessity of tackling these challenges through an integrated, interdisciplinary strategy. We can develop methods for reducing these risks and advancing the preservation of aquatic ecosystems by knowing the historical background of aquatic research threats. In the end, this study shows that maintaining aquatic ecosystems necessitates a consistent dedication to management, conservation, and research. Together, we can tackle the multifaceted risks that aquatic ecosystems face and guarantee the long-term viability and health of these essential ecosystems.

### References

- Appenroth, K.J., 2010.** Definition of “heavy metals” and their role in biological systems. *Soil heavy metals*, pp.19-29. [https://doi.org/10.1007/978-3-642-02436-8\\_2](https://doi.org/10.1007/978-3-642-02436-8_2)
- Bhat, S.U. and Qayoom, U., 2021.** Implications of sewage discharge on freshwater ecosystems.
- Lamberth, C., Jeanmart, S., Luksch, T. and Plant, A., 2013.** Current challenges and trends in the discovery of agrochemicals. *Science*, 341(6147), pp.742-746. <https://doi.org/10.1126/science.1237227>
- Mateo-Sagasta, J., Zadeh, S.M. and Turrall, H. eds., 2018.** More people, more food, worse water?: a global review of water pollution from agriculture.
- Saha, N. and Zaman, M.R., 2011.** Concentration of selected toxic metals in groundwater and some cereals grown in Shibganj area of Chapai Nawabganj, Rajshahi, Bangladesh. *Current Science*, pp.427-431.
- Sullieva, S., Zokirov, K., Nishonov, N., Atamuratova, N. and Abdullayev, B., 2024.** Effect of planting time and seedling feeding area on the yield and biochemical composition of leek (*Allium porrum* L.). In *BIO Web of Conferences* (Vol. 93, p. 02006). EDP Sciences. <https://doi.org/10.1051/bioconf/20249302006>
- SV, B. and Bhandarkar, W.R., 2013.** A study on seasonal variation of physico-chemical properties in some freshwater lotic ecosystems in Gadchiroli District Maharashtra. *Int. J. of Life Sciences*, 1(3), pp.207-215.
- Tiwari, A.K., Singh, A.K., Singh, A.K. and Singh, M.P., 2017.** Hydrogeochemical analysis and evaluation of surface water quality of Pratapgarh district, Uttar Pradesh, India. *Applied Water Science*, 7,

pp.1609-1623.

<https://doi.org/10.1007/s13201-015-0313-z>.

**Vollenweider, S., Grassi, G., König, I. and Puhan, Z., 2003.** Purification and structural characterization of 3-hydroxypropionaldehyde and its derivatives. *Journal of agricultural and food chemistry*, 51(11), pp.3287-3293. <https://doi.org/10.1021/jf021086d>